



POLICY STATEMENT

# The Apgar Score

**American Academy of Pediatrics**

Committee on Fetus and Newborn

**American College of Obstetricians and Gynecologists**

Committee on Obstetric Practice

Organizational Principles to Guide and  
Define the Child Health Care System and/or  
Improve the Health of All Children

## ABSTRACT

The Apgar score provides a convenient shorthand for reporting the status of the newborn infant and the response to resuscitation. The Apgar score has been used inappropriately to predict specific neurologic outcome in the term infant. There are no consistent data on the significance of the Apgar score in preterm infants. The Apgar score has limitations, and it is inappropriate to use it alone to establish the diagnosis of asphyxia. An Apgar score assigned during resuscitation is not equivalent to a score assigned to a spontaneously breathing infant. An expanded Apgar score reporting form will account for concurrent resuscitative interventions and provide information to improve systems of perinatal and neonatal care.

## INTRODUCTION

In 1952, Dr Virginia Apgar devised a scoring system that was a rapid method of assessing the clinical status of the newborn infant at 1 minute of age and the need for prompt intervention to establish breathing.<sup>1</sup> A second report evaluating a larger number of patients was published in 1958.<sup>2</sup> This scoring system provided a standardized assessment for infants after delivery. The Apgar score comprises 5 components: heart rate, respiratory effort, muscle tone, reflex irritability, and color, each of which is given a score of 0, 1, or 2. The score is now reported at 1 and 5 minutes after birth. The Apgar score continues to provide a convenient shorthand for reporting the status of the newborn infant and the response to resuscitation. The Apgar score has been used inappropriately in term infants to predict specific neurologic outcome. Because there are no consistent data on the significance of the Apgar score in preterm infants, in this population the score should not be used for any purpose other than ongoing assessment in the delivery room. The purpose of this statement is to place the Apgar score in its proper perspective.

The neonatal resuscitation program (NRP) guidelines<sup>3</sup> state that “Apgar scores should not be used to dictate appropriate resuscitative actions, nor should interventions for depressed infants be delayed until the 1-minute assessment.” However, an Apgar score that remains 0 beyond 10 minutes of age may be useful in determining whether additional resuscitative efforts are indicated.<sup>4</sup> The current NRP guidelines<sup>3</sup> state that “if there is no heart rate after 10 minutes of complete and adequate resuscitation efforts, and there is no evidence of other causes of newborn compromise, discontinuation of resuscitation efforts may be appropriate. Current data indicate that, after 10 minutes of asystole, newborns are very unlikely to survive, or the rare survivor is likely to survive with severe disability.”

[www.pediatrics.org/cgi/doi/10.1542/peds.2006-0325](http://www.pediatrics.org/cgi/doi/10.1542/peds.2006-0325)

doi:10.1542/peds.2006-0325

All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

### Key Words

Apgar score, asphyxia, neurologic outcome, resuscitation, cerebral palsy

### Abbreviation

NRP—neonatal resuscitation program  
PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275). Copyright © 2006 by the American Academy of Pediatrics and the American College of Obstetricians and Gynecologists

Previously, an Apgar score of 3 or less at 5 minutes was considered an essential requirement for the diagnosis of perinatal asphyxia. *Neonatal Encephalopathy and Cerebral Palsy: Defining the Pathogenesis and Pathophysiology*,<sup>5</sup> produced in 2003 by the American College of Obstetricians and Gynecologists in collaboration with the American Academy of Pediatrics, lists an Apgar score of 0 to 3 beyond 5 minutes as one suggestive criterion for an intrapartum asphyxial insult. However, a persistently low Apgar score alone is not a specific indicator for intrapartum compromise. Further, although the score is used widely in outcome studies, its inappropriate use has led to an erroneous definition of asphyxia. Intrapartum asphyxia implies fetal hypercarbia and hypoxemia, which, if prolonged, will result in metabolic acidemia. Because the intrapartum disruption of uterine or fetal blood flow is rarely, if ever, absolute, asphyxia is an imprecise, general term. Descriptions such as hypercarbia, hypoxia, and metabolic, respiratory, or lactic acidemia are more precise for immediate assessment of the newborn infant and retrospective assessment of intrapartum management.

#### LIMITATIONS OF THE APGAR SCORE

It is important to recognize the limitations of the Apgar score. The Apgar score is an expression of the infant's physiologic condition, has a limited time frame, and includes subjective components. In addition, the biochemical disturbance must be significant before the score is affected. Elements of the score such as tone, color, and reflex irritability partially depend on the physiologic maturity of the infant. The healthy preterm infant with no evidence of asphyxia may receive a low score only because of immaturity.<sup>6</sup> A number of factors may influence an Apgar score, including but not limited to drugs, trauma, congenital anomalies, infections, hypoxia, hypovolemia, and preterm birth.<sup>7</sup> The incidence of low Apgar scores is inversely related to birth weight, and a low score is limited in predicting morbidity or mortality.<sup>8</sup> Accordingly, it is inappropriate to use an Apgar score alone to establish the diagnosis of asphyxia.

#### APGAR SCORE AND RESUSCITATION

The 5-minute Apgar score, and particularly a change in the score between 1 and 5 minutes, is a useful index of the response to resuscitation. If the Apgar score is less than 7 at 5 minutes, the NRP guidelines state that the assessment should be repeated every 5 minutes up to 20 minutes.<sup>3</sup> However, an Apgar score assigned during a resuscitation is not equivalent to a score assigned to a spontaneously breathing infant.<sup>9</sup> There is no accepted standard for reporting an Apgar score in infants undergoing resuscitation after birth, because many of the elements contributing to the score are altered by resuscitation. The concept of an "assisted" score that accounts for

resuscitative interventions has been suggested, but the predictive reliability has not been studied. To describe such infants correctly and provide accurate documentation and data collection, an expanded Apgar score report form is proposed (Fig 1).

#### PREDICTION OF OUTCOME

A low 1-minute Apgar score alone does not correlate with the infant's future outcome. A retrospective analysis concluded that the 5-minute Apgar score remained a valid predictor of neonatal mortality, but using it to predict long-term outcome was inappropriate.<sup>10</sup> On the other hand, another study<sup>11</sup> stated that low Apgar scores at 5 minutes are associated with death or cerebral palsy, and this association increased if both 1- and 5-minute scores were low.

An Apgar score at 5 minutes in term infants correlates poorly with future neurologic outcomes. For example, a score of 0 to 3 at 5 minutes was associated with a slightly increased risk of cerebral palsy compared with higher scores.<sup>12</sup> Conversely, 75% of children with cerebral palsy had normal scores at 5 minutes.<sup>12</sup> In addition, a low 5-minute score in combination with other markers of asphyxia may identify infants at risk of developing seizures (odds ratio: 39; 95% confidence interval: 3.9–392.5).<sup>13</sup> The risk of poor neurologic outcomes increases when the Apgar score is 3 or less at 10, 15, and 20 minutes.<sup>7</sup>

A 5-minute Apgar score of 7 to 10 is considered normal. Scores of 4, 5, and 6 are intermediate and are not markers of increased risk of neurologic dysfunction. Such scores may be the result of physiologic immaturity, maternal medications, the presence of congenital malformations, and other factors. Because of these other conditions, the Apgar score alone cannot be considered evidence or a consequence of asphyxia. Other factors including nonreassuring fetal heart rate monitoring patterns and abnormalities in umbilical arterial blood gases, clinical cerebral function, neuroimaging studies, neonatal electroencephalography, placental pathology, hematologic studies, and multisystem organ dysfunction need to be considered when defining an intrapartum hypoxic-ischemic event as a cause of cerebral palsy.<sup>5</sup>

#### OTHER APPLICATIONS

Monitoring of low Apgar scores from a delivery service can be useful. Individual case reviews can identify needs for focused educational programs and improvement in systems of perinatal care. Analyzing trends allows assessment of the impact of quality improvement interventions.

#### CONCLUSION

The Apgar score describes the condition of the newborn infant immediately after birth<sup>14</sup> and, when properly ap-

**APGAR SCORE**

Gestational Age \_\_\_\_\_ weeks

SIGN	0	1	2	1 minute	5 minute	10 minute	15 minute	20 minute	
				COLOR	Blue or Pale	Acrocyanotic	Completely Pink		
HEART RATE	Absent	<100 minute	>100 minute						
REFLEX IRRITABILITY	No Response	Grimace	Cry or Active Withdrawal						
MUSCLE TONE	Limp	Some Flexion	Active Motion						
RESPIRATION	Absent	Weak Cry Hypoventilation	Good, crying						
TOTAL									
Comments:				Resuscitation					
				Minutes	1	5	10	15	20
				Oxygen					
				PPV/NCPAP					
				ETT					
				Chest Compressions					
				Epinephrine					

FIGURE 1

Expanded Apgar score form. Record the score in the appropriate place at specific time intervals. The additional resuscitative measures (if appropriate) are recorded at the same time that the score is reported using a check mark in the appropriate box. Use the comment box to list other factors including maternal medications and/or the response to resuscitation between the recorded times of scoring. PPV/NCPAP indicates positive-pressure ventilation/nasal continuous positive airway pressure; ETT, endotracheal tube.

plied, is a tool for standardized assessment. It also provides a mechanism to record fetal-to-neonatal transition. An Apgar score of 0 to 3 at 5 minutes may correlate with neonatal mortality but alone does not predict later neurologic dysfunction. The Apgar score is affected by gestational age, maternal medications, resuscitation, and cardiorespiratory and neurologic conditions. Low 1- and 5-minute Apgar scores alone are not conclusive markers of an acute intrapartum hypoxic event. Resuscitative interventions modify the components of the Apgar score. There is a need for perinatal health care professionals to be consistent in assigning an Apgar score during a resuscitation. The American Academy of Pediatrics and the American College of Obstetricians and Gynecologists propose use of an expanded Apgar score reporting form that accounts for concurrent resuscitative interventions.

**AAP COMMITTEE ON FETUS AND NEWBORN, 2005–2006**

- Ann R. Stark, MD, Chairperson
- David H. Adamkin, MD
- Daniel G. Batton, MD
- Edward F. Bell, MD
- Vinod K. Bhutani, MD
- Susan E. Denson, MD
- William A. Engle, MD
- \*Gilbert I. Martin, MD
- Lillian R. Blackmon, MD, Past Chairperson

**LIAISONS**

- Keith J. Barrington, MD  
Canadian Paediatric Society
- Gary D.V. Hankins, MD  
American College of Obstetricians and Gynecologists
- Tonse N.K. Raju, MD, DCH  
National Institutes of Health
- Kay M. Tomashek, MD, MPH  
Centers for Disease Control and Prevention
- Carol Wallman, MSN, RNC, NNP  
National Association of Neonatal Nurses and  
Association of Women’s Health, Obstetric and  
Neonatal Nurses
- Laura E. Riley, MD, Past Liaison  
American College of Obstetricians and Gynecologists

**STAFF**

Jim Couto, MA

**ACOG COMMITTEE ON OBSTETRIC PRACTICE**

- \*Gary D.V. Hankins, MD, Chairperson
- Sarah J. Kilpatrick, MD, Vice-Chairperson
- Angela L. Bell, MD
- Jeanne M. Coulehan, CNM
- Susan Hellerstein, MD
- Jack Ludmir, MD
- Carol A. Major, MD
- Sean McFadden, MD
- Susan M. Ramin, MD
- Russell R. Snyder, Col, MC, USAF

## LIAISONS

Hani K. Atrash, MD, MPH  
Centers for Disease Control and Prevention

William Callaghan, MD  
Centers for Disease Control and Prevention

Joshua A. Copel, MD  
Association for Medical Ultrasound

Gary A. Dildy III, MD  
Society for Maternal-Fetal Medicine

William Herbert, MD  
Committee on Practice Bulletins-Obstetrics Liaison

Samuel C. Hughes, MD  
American Society of Anesthesiologists

Bruce Patsner, MD, JD  
Food and Drug Administration

Colin Pollard  
Food and Drug Administration

Phill Price, MD  
Food and Drug Administration

Catherine Y. Spong, MD  
National Institute of Child Health and Human  
Development

Ann Stark, MD  
American Academy of Pediatrics

John S. Wachtel, MD  
Committee on Quality Improvement and Patient  
Safety

## STAFF

Stanley Zinberg, MD, MS  
Beth Steele  
Debra Hawks, MPH

\*Lead authors

## REFERENCES

1. Apgar V. A proposal for a new method of evaluation of the newborn infant. *Curr Res Anesth Analg*. 1953;32:260–267
2. Apgar V, Holiday DA, James LS, Weisbrot IM, Berrien C. Evaluation of the newborn infant: second report. *JAMA*. 1958;168:1985–1988
3. American Academy of Pediatrics and American Heart Association. *Textbook of Neonatal Resuscitation*. Elk Grove Village, IL: American Academy of Pediatrics and American Heart Association; 2005
4. Jain L, Ferre C, Vidyasagar D, Nath S, Sheftel D. Cardiopulmonary resuscitation of apparently stillborn infants: survival and long-term outcome. *J Pediatr*. 1991;118:778–782
5. American College of Obstetrics and Gynecology, Task Force on Neonatal Encephalopathy and Cerebral Palsy; American Academy of Pediatrics. *Neonatal Encephalopathy and Cerebral Palsy: Defining the Pathogenesis and Pathophysiology*. Washington, DC: American College of Obstetricians and Gynecologists; 2003
6. Catlin EA, Carpenter MW, Brann BS IV, et al. The Apgar score revisited: influence of gestational age. *J Pediatr*. 1986;109:865–868
7. Freeman JM, Nelson KB. Intrapartum asphyxia and cerebral palsy. *Pediatrics*. 1988;82:240–249
8. Hegyi T, Carone T, Anwar M, et al. The Apgar score and its components in the preterm infant. *Pediatrics*. 1998;101:77–81
9. Lopriore E, van Burk F, Walther F, Arnout J. Correct use of the Apgar score for resuscitated and intubated newborn babies: questionnaire study. *BMJ*. 2004;329:143–144
10. Casey BM, McIntire DD, Leveno KJ. The continuing value of the Apgar score for the assessment of the newborn infants. *N Engl J Med*. 2001;344:467–471
11. Moster D, Lie RT, Irgens LM, Bjerkedal T, Markestad T. The association of Apgar score with subsequent death and cerebral palsy: a population-based study in term infants. *J Pediatr*. 2001;138:798–803
12. Nelson KB, Ellenberg JH. Apgar scores as predictors of chronic neurologic disability. *Pediatrics*. 1981;68:36–44
13. Perlman JM, Risser R. Can asphyxiated infants at risk for neonatal seizures be rapidly identified by current high-risk markers? *Pediatrics*. 1996;97:456–462
14. Papile LA. The Apgar score in the 21st century. *N Engl J Med*. 2001;344:519–520

## The Apgar Score

American Academy of Pediatrics, Committee on Fetus and Newborn, American College of Obstetricians and Gynecologists and Committee on Obstetric Practice

*Pediatrics* 2006;117;1444

DOI: 10.1542/peds.2006-0325

### Updated Information & Services

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/117/4/1444>

### References

This article cites 12 articles, 5 of which you can access for free at:  
<http://pediatrics.aappublications.org/content/117/4/1444#BIBL>

### Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):  
**Committee on Fetus & Newborn**  
[http://www.aappublications.org/cgi/collection/committee\\_on\\_fetus\\_newborn](http://www.aappublications.org/cgi/collection/committee_on_fetus_newborn)  
**Fetus/Newborn Infant**  
[http://www.aappublications.org/cgi/collection/fetus:newborn\\_infant\\_sub](http://www.aappublications.org/cgi/collection/fetus:newborn_infant_sub)  
**Neurology**  
[http://www.aappublications.org/cgi/collection/neurology\\_sub](http://www.aappublications.org/cgi/collection/neurology_sub)

### Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

### Reprints

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

# American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **The Apgar Score**

American Academy of Pediatrics, Committee on Fetus and Newborn, American College of Obstetricians and Gynecologists and Committee on Obstetric Practice

*Pediatrics* 2006;117;1444

DOI: 10.1542/peds.2006-0325

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/117/4/1444>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2006 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®

