

Fall Prevention Self-Assessment

Facility Name:

Date:

| Process | Met/Unmet | Notes |
|---|-----------|-------|
| Rounding every 1-2 hours is in place for high-risk patients to address the 3 P's. | | |
| Customized interventions are in place for each patient to reduce the risk of fall and injury. | | |
| Pharmacist is involved in a medication review to avoid unnecessary hypnotics and sedatives. | | |
| A multi-disciplinary team addresses fall prevention strategies. | | |
| The family is involved in efforts to prevent falls. | | |
| Post fall huddles are in place. | | |
| Fall rates are measured and calculated as a rate. | | |
| Fall rates are monitored at least quarterly. | | |
| Fall data is reported on a regular schedule to key stakeholders and staff. | | |
| RCA is conducted for any fall with injury. | | |
| Key Processes in the fall prevention program are monitored for compliance. | | |
| Care plan addressing deficits found on the fall risk assessment has been developed and implemented. | | |
| A method is in place to communicate fall risk from shift to shift among all patient care staff. | | |
| Staff are educated at least annually on patient fall prevention. | | |
| Neurological assessments are conducted on patients who have unwitnessed falls or hit their head. | | |



Fall Prevention Self-Assessment

| Process | Met/Unmet | Notes |
|--|-----------|-------|
| A unit specific PI project is in place for fall prevention to engage staff in fall prevention. | | |
| Fall risk assessment is done upon admission. | | |
| Fall risk is reassessed daily and upon a change in the patient's condition. | | |
| Interventions to prevent hazards of immobility are in place. | | |
| A method is in place to communicate fall risks to all team members. | | |
| No other issues identified. | | |